

Control of *Vibrio alginolyticus* in *Artemia* culture by treatment with bacterial probiotics

L. Villamil, A. Figueras, M. Planas, B. Novoa*

Instituto de Investigaciones Marinas, Consejo Superior de Investigaciones Científicas (CSIC), Eduardo Cabello, 6 Marinas, 36208 Vigo, Spain

Received 8 July 2002; received in revised form 16 September 2002; accepted 16 September 2002

Abstract

In order to determine the influence of six lactic acid bacterial (LAB) strains, lactic acid and major extracellular products (ECPs) of LAB on the bacterial load of *Artemia* culture, two experiments were carried out. In both experiments, the dominant bacterial species were identified as *Vibrio alginolyticus*. The treatment with *Lactobacillus brevis* (1×10^8 bacteria/ml) and lactic acid (0.1 g/l) reduced the load of this bacterium in the *Artemia* culture water.

Biochemical and morphological characteristics of the *V. alginolyticus* strains isolated from *Artemia* culture were determined. For comparative purposes, another *V. alginolyticus* strain isolated from diseased turbot larvae was also included.

The in vitro antibacterial activity of extracellular products (ECPs) from LAB, as well as lactic acid (0.1 and 0.05 g/l), was assessed against both *V. alginolyticus* strains. ECPs from LAB were able to inhibit the growth of *V. alginolyticus*, especially those from *L. brevis*. In the case of lactic acid treatments, no significant inhibition of *V. alginolyticus* growth was observed.

© 2003 Elsevier Science B.V. All rights reserved.

Keywords: *Vibrio alginolyticus*; Lactic acid bacteria; Bacterial probiotics; Lactic acid; Antibacterial activity

1. Introduction

The incidence of microbial diseases has increased dramatically in accordance with the growth of fish larvae production (Toranzo et al., 1994). In the case of turbot

* Corresponding author. Tel.: +34-986-214463; fax: +34-986-292762.

E-mail address: virus@iim.csic.es (B. Novoa).

(*Scophthalmus maximus* L.) culture, high mortalities occur during larval first feeding due to the incidence of pathogenic and opportunistic bacteria (May, 1973; Grisez et al., 1996). Different authors have suggested that this infection happens mainly through the food chain (Sera and Kumata, 1972; Campbell and Buswell, 1983; Muroga et al., 1987) and especially during the feeding with *Artemia* (Chair et al., 1994; Grisez et al., 1996).

One of the most frequently used procedures to avoid the incorporation of undesirable bacteria to the fish culture is the administration of antibiotics in the water (Brown, 1989) or via *Artemia* (Verpraet et al., 1992; Touraki et al., 1999). However, there is a major concern with respect to the use of antibiotics since they could promote the development of antibiotic-resistant bacteria in fish and the environment (Alderman and Hastings, 1998; Brown, 1989; Bjorklund et al., 1991; Smith et al., 1994; Pedersen et al., 1995; Hameed and Balasubramanian, 2000). In order to increase larval survival, several approaches such as improving husbandry nutrition (Reitan et al., 1993; Roennestad et al., 1999), water quality (Vadstein et al., 1993; Skjermo et al., 1997), disinfecting fish eggs (Salvesen and Vadstein, 1995) and UV treatment (Munro et al., 1999) have been proposed. The use of nonspecific immunostimulants (Robertson et al., 1990; Anderson, 1992; Figueras et al., 1997) and bacterial probiotics such as lactic acid bacteria (Gatesoupe, 1991, 1994, 1999; Ringø and Gatesoupe, 1998; Verschuere et al., 2000) have also been proposed.

Probiotics are defined as microbial dietary adjuvants that beneficially affect the host physiology by modulating the mucosal and systemic immunity, as well as improving microbial balance by preventing the colonization of undesirable bacteria in the intestinal tract (Naidu et al., 1999). The role of lactic acid bacteria (LAB) as probiotics in the digestive tract has been extensively studied in endothermic animals and humans (Aiba et al., 1998; Kontula et al., 1998; Kirjavainen et al., 1999). In fish, LAB were recently described as part of the normal microflora (Strøm and Olfasen, 1990; Ringø and Strøm, 1994; Ringø et al., 1998; Robertson et al., 2000). Exogenous LAB administration has been associated with pathogenic bacterial growth inhibition (Lewus et al., 1991; Gildberg et al., 1995; Santos et al., 1996), fish growth promotion (Noh et al., 1994) and in some cases with an increased survival of experimental infected fish (Gatesoupe, 1994; Gildberg et al., 1997; Robertson et al., 2000). Preventing colonization of detrimental bacteria with selected bacterial strains has been proposed as an important alternative for microbial control during *Artemia* culture (Verschuere et al., 1999). It has also been demonstrated that some of these selected bacterial strains can prevent the growth of bacterial pathogens such as *Vibrio proteolyticus* in the *Artemia* culture (Verschuere et al., 2000).

In the current work, we have studied the effect of six LAB strains and lactic acid in the bacterial load of *Artemia* culture. Since we observed an important reduction of *Vibrio alginolyticus* strains (F 236 and F 237) isolated as dominant bacterium in the *Artemia* culture, further experiments were conducted with these isolates and with another *V. alginolyticus* strain (Hq 221) isolated from diseased turbot larvae. The in vitro antibacterial activity of lactic acid and ECPs from LAB against both bacteria was determined. Biochemical and morphological characteristics of all strains were also included.

2. Material and methods

2.1. Lactic acid bacterial strains

Five LAB strains were obtained from the Spanish type culture collection (CECT): *Lactobacillus casei* (CECT 4043), *Lactobacillus brevis* (CECT 815), *Lactobacillus helveticus* (CECT 541), *Lactococcus lactis* spp. *lactis* (CECT 539) and *Leuconostoc mesenteroides* spp. *mesenteroides* (CECT 4046). *Pediococcus acidilactici* (NRRL B-5627) was obtained from the Northern Regional Research Laboratory, USA. All the LAB strains were grown overnight at 30 °C in Man Rogosa and Sharpe broth (MRS), washed and resuspended in sterile phosphate saline buffer (PBS). These LAB species were chosen since they are known to be safe for human use. Furthermore, they were referenced in other investigations as bacterial probiotics in aquaculture or as bacteria with high antimicrobial activity (Schrøde et al., 1980; Shiri Harzevili et al., 1998; Gatesoupe, 1991, 1994, 2002; Nikoskelainen et al., 2001).

2.2. *Artemia* culture

Artemia cysts (INVE) were hatched in filtered seawater at 28–30 °C under continuous aeration and light intensity. After 24 h of incubation, *Artemia* nauplii (instar I) hatched out and were used for experimental studies.

2.3. LAB and lactic acid administration to *Artemia* culture

Six LAB strains were separately administered to *Artemia* nauplii at a final concentration of 1×10^8 bacteria/ml (Vázquez, 2001). In another series of experiments, lactic acid (Sigma) at different final concentrations (0.005, 0.01, 0.05, 0.1 g/l) were administered to *Artemia* nauplii. In both experiments, *Artemia* culture was maintained for 24 h with aeration at 15 °C. After this time, samples of 5 ml were aseptically taken to determine the bacterial load, as described below.

2.4. Sample processing

To separate *Artemia* from culture water, samples from each treatment were passed over a sterile 100- μ m mesh. The *Artemia* trapped in the filter were rinsed with 10 ml of sterile seawater (ssw), resuspended in 1 ml of ssw and aseptically homogenized. Samples (1 ml) from the culture water were aseptically stored in sterile tubes until used.

In order to determine the bacterial load in *Artemia* and culture water, samples were serially diluted and 100 μ l of each dilution was plated on marine agar (MA) (Panreac Quimica) and on thiosulphate citrate bile sucrose (TCBS) (Panreac Quimica). Plates were then incubated for 48–72 h at room temperature (22 °C). After this time, the colonies were counted and the amount of bacteria was calculated as colony forming unit (CFU) per ml. Dominant colony types on plates were visually selected on the basis of different colony appearance and abundance. The selected colonies from each plate were isolated and re-streaked on fresh MA to ensure purity. For long-term preservation,

cultures were frozen at -80°C in tryptone soy broth containing 1% NaCl (TSB-1) and 15% glycerol (v/v).

2.5. Characterization of bacterial isolates

Pure cultures of the bacterial strains were subjected to standard morphological, physiological and biochemical plate and tube tests. Since we focus this study in the *Vibrio* genus, gram character (Buck, 1982), oxidase test, morphology, motility, sensitivity against O/129 and growth on TCBS were the tests used to identify the isolates that are members of this genus. Identification of *V. alginolyticus* species was achieved following the scheme of Alsina and Blanch (1994) and Bergey's Manual of Systematic Bacteriology (1984, vol. 1, section 5). In parallel, the commercial miniaturized Api 20-E (Bio Mérieux) was also used.

Drug resistance patterns of the *V. alginolyticus* isolates were determined by disc diffusion method on Mueller–Hinton agar (Oxoid) supplemented with 1% NaCl. The chemotherapeutic agents and their concentrations ($\mu\text{g}/\text{disc}$) were ampicillin (10), chloramphenicol (30), furazolidone (300), nitrofurantoin (300), oxolinic acid (2), oxytetracycline (30), streptomycin (10), tetracycline (30) and trimethoprim-sulphamethoxazole (25).

The enzymatic activity of the *V. alginolyticus* isolates was determined with the Api ZYM system (Bio Mérieux) according to the manufacturer's instructions. For comparative purposes, *V. alginolyticus* (Hq 221) isolated from diseased turbot larvae was included in all the characterization tests.

2.6. Antibacterial activity of extracellular products from LAB and lactic acid

We determined the in vitro capacity of extracellular products (ECPs) from the six LAB species and lactic acid to inhibit the growth of *V. alginolyticus* strains isolated from *Artemia* culture and from diseased turbot larvae.

2.6.1. Extraction of LAB ECPs

The extraction of ECPs was performed according to Cabo et al. (1999). Briefly, all the LAB strains were grown overnight at 30°C in MRS broth. After incubation, the pH was adjusted to 3.5 with 2 N ClH, then heated at 80°C for 3 min and centrifuged at $500 \times g$ for 30 min. The supernatant containing the LAB ECPs was filtered through a $0.45\text{-}\mu\text{m}$ filter, buffered at pH 6.0 and stored in aliquots at -80°C .

2.6.2. Bactericidal assay

Bacterial strains to be tested were grown overnight at room temperature (22°C) in TSB-1. The assay was performed in triplicate in 96-well plate by dispensing $50\text{ }\mu\text{l}$ of the bacterial suspension (10^8 bacteria/ml) per well and $50\text{ }\mu\text{l}$ of the corresponding treatment: ECPs (undiluted or previously diluted to one half) or lactic acid (0.05 and 0.1 g/l). Controls were included by incubating the bacteria with MRS instead of ECPs, or with PBS at pH 3.5 instead of lactic acid. After 15 h of incubation, changes in optical density (OD)

(600 nm) were measured and the percentage of bacterial survival was calculated as follows:

$$\text{OD treated wells} / \text{OD untreated wells} \times 100.$$

2.7. Statistics

The data were compared using a Student's *t*-test performed using the computer program Excel 98. Results are expressed as the mean \pm standard deviation and the differences were considered significant at $p < 0.05$.

3. Results

3.1. LAB administration to *Artemia* culture

The majority of isolated bacteria from this experiment belonged to the genus *Vibrio*. They were Gram-negative motile rods sensitive to the vibriostatic agent O/129 and were able to grow in TCBS medium. From all the colonies, one with high capacity to lyse surrounding colonies and to overgrow them was isolated and further identified as *V. alginolyticus* (F 236). In the culture water, a total bacterial average of 10^8 CFU/ml was detected (Fig. 1a), while an average of 10^4 CFU/*Artemia* was recorded (Fig. 1b). The treatment with *L. brevis* (CECT 815) and *L. casei* (CECT 4043) almost completely eliminated *V. alginolyticus* from *Artemia*, whereas in water, it was only eliminated by treatment with *L. brevis*.

3.2. Lactic acid administration to *Artemia* culture

Again, in this experiment, most of the isolated bacteria belonged to the *Vibrio* genus. The bacterial colony with high capacity to lyse surrounding colonies and to overgrow them was also identified as *V. alginolyticus* (F 237). The total amount of bacteria isolated from culture water and *Artemia* was similar to that reported in the experiment with LAB treatments. The treatment with the highest concentration of lactic acid (0.1 g/l) diminished the load of *V. alginolyticus* in *Artemia* culture. However, this treatment was not able to completely eliminate this bacterium from *Artemia* culture (Fig. 2).

3.3. Characterization of bacterial isolates

The main biochemical characteristics of *V. alginolyticus* strains isolated from *Artemia* culture (F 236 and F 237) and diseased turbot larvae (Hq 221) are included in Table 1. All bacterial strains presented a similar profile: they were nonpigmented colonies in TSA-1 while in TCBS, they formed yellow (y) colonies. These isolates require Na^+ to grow and were positive for oxidase, catalase, indole and Voges–Proskauer. Their



a.**b.**

Fig. 1. Effect of administration of six LAB strains (1×10^8 bacteria/ml) in the bacterial load of *Artemia* culture. Dominant bacteria isolated from (a) water and from (b) *Artemia* nauplii. Data were expressed as CFU/ml in the case of water and as CFU/*Artemia* in the case of nauplii.

activities with arginine dihydrolase and lysine decarboxylase were negative; however, they differ on ornithine decarboxylase. The strains isolated from *Artemia* culture were positive and the one isolated from the diseased turbot larvae was negative. All isolates presented similar sensitivity patterns with tested antibiotics, chloramphenicol, tetracycline and oxytetracycline being the most effective, while both isolates were resistant to ampicillin (Table 1).

The highest enzymatic activities observed in both *V. alginolyticus* strains were alkaline phosphatase, leucine arylamidase, acid phosphatase and naphthol-AS-bi-phosphohydrolase.




Fig. 2. Effect of administration of lactic acid at different final concentrations (0.005, 0.01, 0.05 and 0.1 g/l) on the bacterial load of *Artemia* culture. Dominant bacteria isolated from (a) water and from (b) *Artemia* nauplii. Data were expressed as CFU/ml in the case of water and as CFU/*Artemia* in the case of nauplii.

The bacterial strains isolated from *Artemia* culture (F 236 and F 237) exhibited a strong activity of α -chymotrypsin, whereas the strain isolated from diseased turbot larvae (Hq 221) showed a weak reaction.

3.4. Bactericidal assay

V. alginolyticus isolated from *Artemia* culture and diseased turbot larvae were sensitive to ECPs from some LAB strains. ECPs from *L. brevis* caused the strongest

Table 1

Main phenotypic characteristics of the strains of *V. alginolyticus* isolated from diseased turbot larvae (Hq 221) and from *Artemia* culture (F 236–F 237)

Characteristics	<i>V. alginolyticus</i> (F 236 and F 337)	<i>V. alginolyticus</i> (Hq 221)
Gram	—	—
Motility	+	+
Oxidase	+	+
Catalase	+	+
Pigment	—	—
TCBS	+	+
Swarming on solid media	+	+
Arginine dihydrolase	—	—
Ornithine decarboxylase	+	—
Lysine decarboxylase	+	+
O/F glucose	+/+	+/+
Citrate	+	+
Indole	+	+
Voges–Proskauer	+	+
ONPG	—	—
H ₂ S production	—	—
Na ⁺ required for growth	+	+
Gelatinase	+	+
Urease	—	—
<i>Growth at (C):</i>		
4	—	—
15	+	+
37	+	+
<i>Acid from:</i>		
Glucose	+	+
Mannitol	+	+
Inositol	—	—
Sorbitol	—	—
Rhamnose	—	—
Sucrose	+	+
Melibiose	—	—
Amygdalin	—	—
<i>Resistance/sensitivity to:</i>		
O/129	S	S
Ampicillin	R	R
Chloramphenicol	S	S
Nitrofurantoin	S	S
Oxolinic acid	S	S
Oxytetracycline	S	S
Streptomycin	S	S
Tetracycline	S	S
Trimethoprim-sulphamethoxazole	S	S

R: resistant; S: sensitive.

inhibition of both *V. alginolyticus* strains, even if diluted to one half. However, diluted ECPs from other LAB were not able to cause a significant reduction in the growth of *V. alginolyticus* (Fig. 3).

In the case of lactic acid, a slight but not significant inhibition of *V. alginolyticus* growth was observed with the highest dose used (0.1 g/l) (data not shown). Higher lactic acid concentrations were not assessed since they were found to be toxic for *Artemia* culture.




Fig. 3. Growth inhibition of *V. alginolyticus* isolated from (a) *Artemia* culture and from (b) diseased turbot larvae by incubation with ECPs (direct or previously diluted to 1/2) from LAB for 15 h. Data were presented as mean percentage of bacterial survival \pm SD.

4. Discussion

Bacteria from the *Vibrio* genus are most frequently associated with mortalities in marine fish farms (Horne et al., 1977; Devesa et al., 1985; Toranzo et al., 1993). *V. alginolyticus* has been described as a pathogenic bacteria for turbot (Austin et al., 1993), sea bream (*Sparus aurata* L.), (Colomi et al., 1981; Balebona et al., 1998) and grouper (*Epinephelus malabaricus*) (Lee, 1995). It has also been reported as opportunistic bacteria for sea mullet (*Mugil cephalus* L.) (Burke and Rodgers, 1981) and red sea bream (*Pagrus major*) (Muroga et al., 1987), while other studies described this bacteria as common intestinal microflora in several marine fish species (Grisez et al., 1997).

Manipulation of bacterial load present in rotifers and *Artemia* for fish feeding may constitute a valuable mechanism to increase larval growth and survival rates (Noh et al., 1994; Gatesoupe, 1994; Robertson et al., 2000). Results obtained in the present study suggest that treatments with *L. brevis* and *L. casei* significantly diminish *V. alginolyticus* load in *Artemia* culture. In the case of lactic acid, only the highest dose used was able to reduce bacterial numbers mainly in the culture water. During the metabolism, LAB produce a range of ECPs that could inhibit bacterial growth such as lactic acid (Lindgren and Clevström, 1978; Alakomi et al., 2000), organic acids (Midolo et al., 1995), hydrogen peroxide, carbon dioxide (Naidu et al., 1999) and bacteriocins. Bacteriocins are bactericidal or bacteriostatic polypeptides that are mostly active against closely related bacteria (Klaenhammer, 1998) and gram-positive microbes (Galvin et al., 1999; Hurst, 1981; Severina et al., 1998). However, some LAB also seem to be antagonistic against gram-negative bacteria such as *Vibrio anguillarum*, *Vibrio salmonicida* and *Proteus vulgaris* (Strøm, 1998 in Ringø and Gatesoupe, 1988). The carnocin from *Carnobacterium piscicola* was also found to be effective against *Aeromonas hydrophila* (Lewus et al., 1991).

In vitro bactericidal activity results were well correlated with in vivo observations. Experiments showed that lactic acid slightly, but not significantly, inhibited *V. alginolyticus* growth; however, a major growth inhibition was obtained when the bacteria were incubated with ECPs from LAB. These results indicate that the bacterial inhibition may be caused by the synergism of various compounds produced as ECPs of LAB and not by lactic acid alone. Undissociated weak acids like lactic acid possess the ability to cross membranes of microorganisms, become dissociated inside and acidify the interior causing the expulsion of H^+ ions from the cells; this action weakens the cells and makes them more susceptible to bacteriocins and other bactericidal compounds (Nykänen et al., 1998). Other mechanisms that could inhibit the growth of undesirable bacteria such as competition for nutrients, available energy or adhesion sites (Gatesoupe, 1999; Verschuere et al., 2000) should be considered. LAB have also been successfully applied in the rotifer (*Brachionus plicatilis*) culture. Gatesoupe (1991) reported that a commercial preparation of live *Lactobacillus plantarum* decreased the proportion of *Aeromonas salmonicida*-like bacteria associated with rotifers, and Harzevili et al. (1988) demonstrated that *L. lactis* (AR21) exhibited an inhibitory effect against *V. anguillarum* in rotifer culture at suboptimal conditions.

Treatment of *Artemia* with acids such as hypochlorite has been demonstrated to completely disinfect cysts. However, these are rapidly recolonized with bacteria (Sogerloos et al., 2001). In order to maintain the inhibition of undesirable bacteria, it is possible

that LAB, as well as chemical compounds, have to be supplied on a regular basis. Jöborn et al. (1997) reported that the addition of high doses of LAB caused a temporary change in the composition of fish intestinal microbial communities, but within a few days after the intake had stopped, the exogenous added bacteria showed decrease in numbers and were lost from the gastrointestinal tract of most of the fish.

Further studies, considering growth and survival capacity at longer time periods, will clarify the effectiveness of LAB as biological control for undesirable bacteria during turbot culture.

Acknowledgements

This work was partially supported by the projects 1FD97-0044-C03-03 and 1FD97-0044-C03-02 from FEDER and a grant from Caixa Galicia (Spain). We would like to thank Dr. Murado for the supply of LAB for in vivo experimental assays, Stolt Sea Farm and specially Ana Riaza for their support. L. Villamil acknowledges the University of Vigo for a research fellowship.

References

- Aiba, Y.S., Nobuyuki, M., Abu, M., Kabir, T., Atushi, Yasuhiro Koga, M.Y., 1998. Lactic acid-mediated suppression of *Helicobacter pylori* by the oral administration of *Lactobacillus salivarius* as a probiotic in a gnotobiotic murine model. *Am. J. Gastroenterol.* 93, 2097–2101.
- Alakomi, H.L., Skyttä, E., Saarela, M., Mattila-Sandholm, T., Latva-kala, K., Helander, I.M., 2000. Lactic acid permeabilizes gram-negative bacteria by disrupting the outer membrane. *Appl. Environ. Microbiol.* 66, 2001–2005.
- Alderman, D.J., Hastings, T.S., 1998. Antibiotic use in aquaculture: development of antibiotic resistance—potential for consumer risks. *Int. J. Food Sci. Technol.* 33, 139–155.
- Alsina, M., Blanch, A., 1994. Improvement and update of a set of keys for biochemical identification of *Vibrio* species. *J. Appl. Bacteriol.* 77, 719–721.
- Anderson, D.P., 1992. Immunostimulants, adjuvants and vaccine carriers in fish: applications to aquaculture. *Ann. Rev. J. Fish Dis.* 2, 281–307.
- Austin, B., Stobie, M., Robertson, P.A.W., Glass, H.S., Mudarris, M., Stark, J.R., 1993. *Vibrio alginolyticus*: the cause of gill disease leading to progressive low-level mortalities among juvenile turbot, *Scophthalmus maximus* L., in a Scottish aquarium. *J. Fish Dis.* 16, 277–280.
- Balebona, M.C., Andreu, M.J., Bordas, M.A., Zorrilla, I., Moriñigo, M.A., Borrego, J.J., 1998. Pathogenicity of *Vibrio alginolyticus* for cultured gilt-head sea bream (*Sparus aurata* L.). *Appl. Environ. Microbiol.* 64, 4269–4275.
- Bergey's Manual of Systematic Bacteriology, vol. 1. Williams & Wilkins, Baltimore, USA. 964 pp.
- Bjorklund, H.V., Rabeg, C.M.I., Bylund, G., 1991. Residues of oxolinic acid and oxytetracycline in fish and sediments from fish farms. *Aquaculture* 97, 85–96.
- Brown, J.H., 1989. Antibiotics: their use and abuse in aquaculture. *World Aquac.* 20, 34–42.
- Buck, I.D., 1982. Non staining (KOH) method for determination of gram reactions of marine bacteria. *Appl. Environ. Microbiol.* 44, 992–993.
- Burke, J., Rodgers, L., 1981. Identification of pathogenic bacteria associated with the occurrence of “red spot” in sea mullet, *Mugil cephalus* L., in south-eastern Queensland. *J. Fish Dis.* 3, 153–159.
- Cabo, M.L., Murado, M.A., González, M.P., Pastoriza, L., 1999. A method for bacteriocin quantification. *J. Appl. Microbiol.* 87, 907–914.
- Campbell, A.C., Buswell, J.A., 1983. The intestinal microflora of farmed Dover sole (*Solea solea*) at different stages of fish development. *J. Appl. Bacteriol.* 35, 215–223.

- Chair, M., Dehasque, M., Van Pouke, S., Nelis, H., Sogerloos, P., De Leenheer, A.P., 1994. An oral challenge for turbot larvae with *Vibrio anguillarum*. *Aquac. Int.* 2, 270–272.
- Colomi, A., Paperna, I., Gording, H., 1981. Bacterial infections in gilt-head sea bream *Sparus aurata* cultured at Elat. *Aquaculture* 23, 257–267.
- Devesa, S., Toranzo, A.E., Barja, J.L., 1985. First report of vibriosis in turbot (*Scophthalmus maximus*) cultures in northwestern Spain. In: Ellis, A.E. (Ed.), *Fish Shellfish Path.* Academic Press, London, pp. 131–140.
- Figueras, A., Santarem, M., Novoa, B., 1997. In vitro immunostimulation of turbot (*Scophthalmus maximus*) leukocytes with β glucan and/or *Photobacterium damsela* bacterin. *Fish Pathol.* 32, 153–159.
- Galvin, M., Hill, C., Ross, R.P., 1999. Lacticin 3147 displays activity in buffer against gram-positive bacterial pathogens, which appear insensitive in standard plate assays. *Lett. Appl. Microbiol.* 28, 355–358.
- Gatesoupe, F.J., 1991. The effect of three strains of lactic acid bacteria on the production rate of rotifers *Brachionus plicatilis*, and their dietary value for larval turbot *Scophthalmus maximus*. *Aquaculture* 96, 335–342.
- Gatesoupe, F.J., 1994. Lactic acid bacteria increase the resistance of turbot larvae, (*Scophthalmus maximus*) against pathogenic vibrio. *Aquat. Living Resour.* 7, 182–277.
- Gatesoupe, F.J., 1999. The use of probiotics in aquaculture. *Aquaculture* 180, 147–165.
- Gatesoupe, F.J., 2002. Probiotic and formaldehyde treatment of *Artemia* nauplii as food for larval pollack, *Pollachius pollachius*. *Aquaculture* 212, 347–360.
- Gildberg, A., Johansen, A., Bøgwald, J., 1995. Growth and survival of Atlantic salmon (*Salmo salar*) fry given diets supplements with fish protein hydrolysate and lactic acid bacteria during a challenge trial with *Aeromonas salmonicida*. *Aquaculture* 138, 23–34.
- Gildberg, A., Mikkelsen, H., Sandaker, E., Ringø, E., 1997. Probiotic effect of lactic acid bacteria in the feed on growth and survival of fry of Atlantic cod (*Gadus morhua*). *Hydrobiologia* 352, 279–285.
- Grisez, L., Chair, M., Sogerloos, P., Ollevier, F., 1996. Mode of infection and spread of *Vibrio anguillarum* in turbot *Scophthalmus maximus* larvae after oral challenge through live feed. *Dis. Aquat. Org.* 26, 181–187.
- Grisez, L., Reyniers, J., Verdonck, L., Swings, J., Ollevier, F., 1997. Dominant intestinal microflora of sea bream and sea bass larvae, from two hatcheries, during larval development. *Aquaculture* 155, 377–399.
- Hameed, A.S., Balasubramanian, G., 2000. Antibiotic resistance in bacteria isolated from *Artemia* nauplii and efficacy of formaldehyde to control bacterial load. *Aquaculture* 183, 195–205.
- Harzevili, A.R.S., Van Duffel, H., Dhert, P., Swings, J., Sogerloos, P., 1998. Use of a potential probiotic *Lactococcus lactis* AR21 strain for the enhancement of growth in the rotifer *Brachionus plicatilis* (Müller). *Aquac. Res.* 29, 411–417.
- Horne, M.T., Richards, R.H., Roberts, R.J., Smith, P.C., 1977. Peracute vibriosis in juvenile turbot *Scophthalmus maximus*. *J. Fish Biol.* 11, 355–361.
- Hurst, A., 1981. Nisin. *Adv. Appl. Microbiol.* 27, 85–123.
- Jöborn, A., Olsson, J.C., Westerdlh, P., Conway, L., Kjelleberg, S., 1997. Colonization in the fish intestinal tract and production of inhibitory substances in intestinal mucus and fecal extracts by *Carnobacterium* sp. strain K1. *J. Fish Dis.* 20, 383–392.
- Kirjavainen, P.V., El-Nezami, H.S., Salminen, S.J., Ahokas, J.T., Wright, P.F.A., 1999. The effect of orally administrated viable probiotic and dairy lactobacilli on mouse lymphocyte proliferation. *FEMS Immunol. Med. Microbiol.* 26, 131–135.
- Klaenhammer, T.R., 1998. Bacteriocins of lactic acid bacteria. *Biochimie* 70, 337–349.
- Kontula, P., Jaskari, J., Nollet, L., De Smet, I., Von Wright, A., Poutanen, K., Mattila-Sandholm, T., 1998. The colonization of a simulator of the human intestinal microbial ecosystem by a probiotic strain fed on a fermented oat bran product: effects on the gastrointestinal microbiota. *Appl. Microbiol. Biotechnol.* 50, 246–252.
- Lee, K.K., 1995. Pathogenesis studies on *Vibrio alginolyticus* in the grouper, *Epinephelus malabaricus*, Bloch et Schneider. *Microb. Pathog.* 19, 39–48.
- Lewus, C.B., Kaiser, A., Montville, T.J., 1991. Inhibition of food-borne pathogens by bacteriocins from lactic acid bacteria isolated from meat. *Microbiology* 57, 1683–1688.
- Lindgren, S., Clevström, G., 1978. Antibacterial activity of lactic acid bacteria. *Swed. J. Agric. Res.* 8, 61–66.
- May, R.C., 1973. Larval mortality in marine fishes and the critical period concept. In: Blaxter, J.H.S. (Ed.), *The Early Life History of Fish*. Springer-Verlag, Berlin, pp. 3–19.

- Midolo, P.D., Lambert, J.R., Hull, R., Luo, F., Grayson, M.L., 1995. In vitro inhibition of *Helicobacter pylori* NCNT 11637 by organic acids and lactic acid bacteria. J. Appl. Bacteriol. 79, 475–479.
- Munro, P.D., Henderson, P.J., Barbour, A., Birkbeck, T.H., 1999. Partial decontamination of rotifers with ultra-violet radiation: the effect of change in the bacterial load and flora of rotifers on mortalities in start-feeding larval turbot. Aquaculture 170, 229–244.
- Muroga, K., Higashi, M., Keetoku, H., 1987. The isolation of intestinal microflora of farmed red seabream (*Pagrus major*) and black seabream (*Acanthopagrus schlegelii*) at larval and juvenile stages. Aquaculture 65, 79–88.
- Naidu, A.S., Bidlack, W.R., Clemens, R.A., 1999. Probiotic spectra of lactic acid bacteria. Crit. Rev. Food Sci. 39, 13–126.
- Nikoskelainen, S., Salminen, S., Bylund, G., Ouwehand, A.C., 2001. Characterization of the properties of human- and dairy-derived probiotics for prevention of infectious diseases in fish. Appl. Environ. Microbiol. 67, 2430–2435.
- Noh, S.H., Han, K., Won, T.H., Choi, Y.J., 1994. Effect of antibiotics, enzyme, yeast culture and probiotics on the growth performance of Israeli carp. Korean J. Anim. Sci. 36, 480–486.
- Nykänen, A., Vesanen, S., Kallio, H., 1998. Synergistic antimicrobial effect of nisin whey permeate and lactic acid in microbes isolated from fish. Lett. Appl. Microbiol. 27, 345–348.
- Pedersen, K., Tiainen, T., Larsen, J.L., 1995. Antibiotic resistance of *Vibrio anguillarum*, in relation to serovar and plasmid contents. Acta Vet. Scand. 36, 55–64.
- Reitan, K.I., Rainuzzo, J.R., Oie, G., Olsen, Y., 1993. Nutritional effects of algal addition in first-feeding of turbot (*Scophthalmus maximus*) larvae. Aquaculture 118, 257–275.
- Ringø, E., Gatesoupe, F.J., 1998. Lactic acid bacteria in fish: a review. Aquaculture 160, 177–203.
- Ringø, E., Strøm, E., 1994. Microflora of Atlantic char, *Salvelinus alpinus* (L.); gastrointestinal flora microflora of free-living fish, and effect of diet and salinity on the intestinal microflora. Aquac. Fish. Manage. 25, 623–629.
- Ringø, E., Bendisken, H.R., Gausen, S.J., Sundsfjord, A., Olsen, R.E., 1998. The effect of dietary fatty acids on lactic acid bacteria associated with the epithelial mucosa and from faecalia of Atlantic charr, *Salvelinus alpinus* (L.). J. Appl. Microbiol. 85, 855–864.
- Robertsen, B., Rørstad, G., Engstad, R., Raa, J., 1990. Enhancement of non-specific disease resistance in Atlantic salmon, *Salmo salar* L., by a glucan from *Saccharomyces cerevisiae* cell wall. J. Fish Dis. 13, 391–400.
- Robertson, P.A.W., O'Dowd, C., Burrells, C., Williams, P., Austin, B., 2000. Use of *Carnobacterium* sp. a probiotic for Atlantic salmon (*Salmo salar* L.) and rainbow trout (*Oncorhynchus mykiss*, Walbaum). Aquaculture 185, 235–243.
- Roennestad, I., Thorsen, A., Finn, R.N., 1999. Fish larval nutrition: a review of recent advances in the roles of amino acids. Aquaculture 177, 1201–1216.
- Salvesen, I., Vadstein, O., 1995. Surface disinfection of eggs from marine fish: evaluation of four chemicals. Aquac. Int. 3, 155–177.
- Santos, J., Lopez-Díaz, T., Garcia-Fernandez, M., Garcia-Otero, A., 1996. Effect of lactic starter culture on the growth and protease activity of *Aeromonas hydrophila*. J. Appl. Bacteriol. 80, 13–18.
- Schrøde, K., Clausen, E., Sandberg, A.M., Raa, J., 1980. Psychrotrophic *Lactobacillus plantarum* from fish and its ability to produce antibiotic substances. In: Connell, J.J. (Ed.), Advances in Fish Science and Technology. Fishing News Books, Farnham, Surrey, England, pp. 480–483.
- Sera, H., Kumata, M., 1972. Bacterial flora in the digestive tract of marine fish. Bacterial flora of fish, red seabream snapper and crimson sea bream, fed three kind of diets. Bull. Jap. Soc. Sci. Fish. 38, 50–55.
- Severina, E., Severin, A., Tomasz, A., 1998. Antibacterial efficacy of nisin against multidrug-resistant gram-positive pathogens. J. Antimicrob. Chemother. 4, 341–347.
- Shiri Harzevili, A.R., Van Duffel, H., Dhert, P., Swings, J., Sogerloos, P., 1998. Use of a potential probiotic *Lactococcus lactis* AR21 strain for the enhancement of growth in the rotifer *Brachionus plicatilis* (Müller). Aquac. Res. 29, 411–417.
- Skjermo, J., Salvesen, I., Oie, G., Olsen, Y., Vadstein, O., 1997. Microbially matured water: a technique for selection of a non-opportunistic bacterial flora in water that may improve performance of marine larvae. Aquac. Int. 5, 13–28.
- Smith, P., Hiney, M.P., Samuelsen, B.D., 1994. Bacterial resistance to antimicrobial agents used in fish farming: a critical evaluation of method and meaning. Ann. Rev. J. Fish Dis. 4, 273–313.

- Sogerloos, P., Dhert, P., Candevra, P., 2001. Use of brine shrimp, *Artemia* spp., in marine fish larviculture. *Aquaculture* 200, 147–159.
- Strøm, E., Olfasen, J.A., 1990. The indigenous flora of wild-captured juvenile cod in net-pen rearing. In: Lèsel, R. (Ed.), *Microbiology in Poecilotherms*. Elsevier, Amsterdam, pp. 181–185.
- Toranzo, A.E., Novoa, B., Romalde, J.L., Núñez, S., Devesa, S., Mariño, E., Silva, R., Martínez, E., Figueras, A., Barja, J.L., 1993. Microflora associated with healthy and diseased turbot (*Scophthalmus maximus*) from three farms in northwest Spain. *Aquaculture* 114, 189–202.
- Toranzo, A.E., Barja, J.L., Devesa, S., 1994. An overview of the main infectious problems in cultured turbot: present status and future necessities. *Eur. Aquac. Soc., Spec. Publ.* 22, 106–126.
- Touraki, M., Niopas, I., Kastritsis, C., 1999. Bioaccumulation of trimethoprim, sulfamethoxazole and *N*-acetyl-sulfamethoxazole in *Artemia* nauplii and residual kinetics in seabass larvae after repeated oral dosing of medicated nauplii. *Aquaculture* 175, 15–30.
- Vadstein, O., Oie, G., Salvesen, I., Skjermo, J., Skjak-Braek, G., 1993. A strategy to obtain microbial control during larval development of marine fish. In: Reirtesen, Dahle, Jorgesen, Tvinnereim (Eds.), *Balkema*, Rotterdam, pp. 69–75.
- Vázquez, J.A., 2001. Lactobacterias como probiontes e produtoras de bacteriocinas. Modelos de crecimiento e actividade. Aplicacións á acuicultura, PhD thesis. University of Santiago de Compostela (In Gallego). 346 pp.
- Verpraet, R., Chair, M., Léger, P., Nelis, H., Sogerloos, P., De Leenheer, A., 1992. Live-food mediated drug delivery as a tool for disease treatment in larviculture. The enrichment of therapeutics in rotifers and *Artemia* nauplii. *Aquac. Eng.* 11, 113–139.
- Verschuere, L., Rombaut, G., Huys, G., Dhont, J., Sogerloos, P., Verstrate, W., 1999. Microbial control through pre-emptive colonization by selected bacterial strains. *Appl. Environ. Microbiol.* 65, 2527–2533.
- Verschuere, L., Heang, H., Criel, G.R., Sogerloos, P., Verstrate, W., 2000. Selected bacterial strains protect *Artemia* spp. from the pathogenic effects of *Vibrio proteolyticus* CW8T2. *Appl. Environ. Microbiol.* 66, 1139–1146.